

Assessment and Diagnosis of Substance Use Disorders

Ken Weingardt, Ph.D.
Center for Health Care Evaluation
VA Palo Alto Health Care System and
Stanford University School of Medicine
Palo Alto, CA









Detection of potential problem






















Screening



<u>Definition</u>: "The skillful use of empirically based procedures for identifying individuals who have SUDs or who are at risk for developing them".


Primary Goal: Detect individuals with SUDs

Modalities:

Biological markers (GGT, MCV) Self-report measures



Screening



Result of Screening Measure


| SUD | Negative | Positive |
|---------|--------------------|--------------------|
| Absent | True Negative (a) | False Positive (b) |
| Present | False Negative (c) | True Positive (d) |

<u>Sensitivity</u>: proportion of persons with SUD that are correctly (true positive) by the test (d/d+c).

Specificity: proportion of persons without SUD Correctly identified as true negatives (a/a+b)



Screening



Guidelines for selecting screening measures

- The goals of the screening
- 2. The time available for conducting the screening
- 3. The resources available for scoring the screening test and providing feedback/referral for positive Cases.

Comparison among screening measures



CAGE




MAST



AUDI7








What are the parameters of the target problem?

- 1. Measures of consumption Quantity and frequency
- 2. Measures of consequences

 Areas of impact of SUD in other areas of life
- 3. Comprehensive measures
 - o Family/Social
 - o Psychiatric
 - o Medical
 - o Legal
 - o Employment







1. Quantity and frequency of consumption


<u>Summary/aggregate measures</u> of consumption: Respondents summarize recent use patterns *Number of days in the past 30 that you:*

<u>Daily diary/daily drinking/daily use measures</u>: Respondents report all of their recent use/drinking *Timeline Follow-back*

Primary versus Composite Measures








2. Measures of consequences

Drinkers Inventory of Consequences (DrInC)
Measures adverse consequences of alcohol use:

- Interpersonal
- Physical
- Social
- Impulsive
- Intrapersonal







3. Comprehensive Measures

Addiction Severity Index



Semi-structured interview designed to address seven Potential problem areas in substance abusing patients


- Medical status
- Employment & Support
- Drug use
- Alcohol use
- Legal status
- Family/social status
- Psychiatric status





"Get the ASI-MV and all the uses you need via phone, fax or email"





Does problem meet diagnostic criteria?

Substance Abuse


A maladaptive pattern of substance use leading to clinically significant impairment or distress, as manifested by one (or more) of the following occurring within a 12 month period:

- Recurrent substance use resulting in a failure to fulfill major role obligations at work, school or home.
- Recurrent substance use in situation in which it is physically hazardous (e.g. DWI, operating machinery)
- Recurrent substance-related legal problems
- Continued substance use despite having persistent or recurrent social or interpersonal problems caused or exacerbated by the effects of the substance (e.g. arguments with spouse about consequences of intoxication, physical fights)

B. The symptoms have never met the criteria for Substance Dependence for this class of substance.







Does problem meet diagnostic criteria?


Substance Dependence

A maladaptive pattern of substance use, leading to clinically significant impairment or distress, as manifested by three (or more) of the following, occurring at any time during the same 12 month period.

- (1) Tolerance, as defined by either of the following
 - (a) A need for markedly increased amounts of the substance to achieve intoxication or desired effect
 - (b) Markedly diminished effect with continued use of the same amount of the substance
- (2) Withdrawal, as defined by either of the following:
 - (a) the characteristic withdrawal syndrome for the substance
 - (b) the same (or a closely related) substance is taken to relieve or avoid withdrawal symptoms.








Does problem meet diagnostic criteria?

Substance Dependence continued

- (3) The substance is taken in larger amounts or over a longer period than was intended
- (4) There is a persistent desire or unsuccessful efforts to cut down or control substance use
- (5) A great deal of time is spent in activities necessary to obtain the substance (e.g. visiting multiple doctors), use the substance & recover from its effects.
- (6) Important social, occupational or recreational activities are given up or reduced because of substance use
- (7) The substance use is continued despite knowledge of having a persistent or recurrent physical or psychological problem that is likely to have been caused or exacerbated by the substance.



- Brief intervention
- Appropriate referrals
- Specialty psychosocial treatment
- Pharmacotherapy
- Care management



And now for a word from our sponsor.... www.detoxguideline.org



VA/DoD Clinical Practice Guidelines



Management of Alcohol Withdrawal

Course Map

The map below indicates your progress through the course. You may navigate to a module by clicking on its corresponding symbol. You may access the Course Map link located on the left navigation bar at any time during the course.

Home

Getting Started

About

Feedback

Downloads

Help

Quick Reference

Acknowledgements

